MASS LUNG & Allergy PC

Medications That Affect Allergy Skin Testing

Below are medications that MUST be stopped a minimum of 7-10 days prior to allergy testing

Allergy Medications:

Allegra	(Generic: Fexofenadine)
Atarax	(Generic: Hydroxyzine)
Benadryl	(Generic: Diphenhydramine
Chlor-Trimeton	(Generic: Chlorpheniramine)
Astelin/Astepro	Nasal Sprays

Generic: Loratadine)
(Generic: Desloratadine)
(Generic: Levocetirizine)
(Generic: Cetirizine)
Eye Drops

The following heart burn medications also **DO AFFECT** allergy testing:

Pepcid	(Generic: Famotidine)	Tagamet	(Generic: Cimetidine)
Zantac	(Generic: Ranitidine)	Tylenol PM	(And <u>all</u> other "PM" medications)

Please contact the office if you are taking any of the following Medications:

Elavil	(Generic: Amitriptyline)	
Kenalog	(Generic: Triamcinolone)	
Medrol	(Generic: Methylprednisolone)	
Nortriptyline	a	

Prednisone Sinequan Seroquel Trazodone

(Generic: **Doxepin**) (Generic: **Quetiapine**)

If you have any questions about your medications, please call the office:

(774) 420-2611